



FACTORS INFLUENCING CONVERSION FROM LAPAROSCOPIC TO OPEN CHOLECYSTECTOMY DUE TO RISK OF BILE DUCT INJURY: AN ANALYTICAL STUDY

Shahnawaz Abro¹, Shahida Baloch², Sajjad Hussain Qureshi³, Syed Kashif Ali Shah⁴, Naeem Al Karim Bhatti⁵, Fouzia Nusrat Pathan⁶.

ABSTRACT

BACKGROUND: Cholecystectomy is now one of the most commonly performed surgical procedures because the incidence of gallbladder disease has risen dramatically. The latter that is, LC is preferred than OC since it is less invasive, causes less pain and has shorter hospital stay. However, LC has been attributes to pose a threat to the BDI and as a result requires conversion to OC and which in the process prolongs the operating time, hospital stay and consequently adds more morbidity. **OBJECTIVE:** This study aims to identify and analyze factors influencing the conversion from laparoscopic cholecystectomy to open cholecystectomy due to the risk of bile duct injury. **MATERIALS AND METHODS:** This was a six months prospective observational study conducted on surgical units I, II, III of Peoples Medical College Hospital, Nawabshah. We included CHC in a total of 132 patients undergoing elective or emergency cholecystectomy who had no previous upper abdominal surgery, liver or bile duct disease not attributable to gallbladder disease, and impaired consent. Data collection included baseline demographic data, clinical presentation, imaging data, intraoperative type of procedure, complications, conversion reasons, and a postoperative complications, recovery outcome data. We performed statistical analysis using SPSS version 25, and logistic regression to determine the significant predictors of conversion. **RESULTS:** Of the 132 patients mean age 45.6 ± 12.3 years, 56.1% were female. Comorbidities included hypertension 24.2% and diabetes 20.5%. Abdominal pain was the most common preoperative symptom 72%. Conversion from LC to OC occurred in 22 out of 112 LCs 19.6%, primarily due to difficult anatomy 45.5%, risk of BDI 31.8%, severe inflammation/adhesions 18.2%, and intraoperative bleeding 4.5%. Logistic regression analysis revealed that age, comorbidities especially hypertension, difficult anatomy, and abnormal preoperative imaging were significant predictors of conversion $p < 0.05$, while gender was not. **CONCLUSION:** This study has shown that challenging anatomic features and patient-specific factors like age and comorbidities are valuable predictive factors for conversion from LC to OC because of BDI concerns. This study suggests that enhanced preoperative evaluation of the patients and isolation of the presumably risky ones will help decrease the rates of the conversion and the possible adverse consequences.

KEYWORDS: Laparoscopic cholecystectomy, open cholecystectomy, bile duct injury, conversion, difficult anatomy, comorbidities, surgical outcomes.

1. MBBS, MS General Surgery Associate Professor Indus Medical College Tando Muhammad Khan.
2. MBBS, MS General Surgery Assistant Professor Department of Surgery Peoples University of Medical & Health Sciences for Women PUMHSW Nawabshah SBA.
3. MBBS, MS General Surgery Assistant Professor Department of Surgery Peoples University of Medical & Health Sciences for Women PUMHSW Nawabshah SBA.
4. MBBS, MS General surgery Associate Professor Department of Surgery Peoples University of Medical & Health Sciences for Women PUMHSW Nawabshah SBA.
5. Mbbs, Ms general surgery Assistant Professor Department Of Surgery Peoples University Of Medical & Health Sciences For Women Pumhsw Nawabshah SBA.
6. Associate Professor, OBS/GYN, KMC Khairpur.

Corresponding Author: Dr Sajjad Hussain Qureshi MBBS, MS General Surgery Assistant Professor Department of Surgery Peoples University of Medical & Health Sciences for Women PUMHSW Nawabshah SBA Email: farooq346@gmail.com. Cell:03347196442

How to Cite This Article: Abro S¹, Baloch S², Qureshi SH³, Shah SKA⁴, Bhatti NAK⁵, Pathan NF⁶ **FACTORS INFLUENCING CONVERSION FROM LAPAROSCOPIC TO OPEN CHOLECYSTECTOMY DUE TO RISK OF BILE DUCT INJURY: AN ANALYTICAL STUDY.** JPUMHS;2024;14:03,103-111. <http://doi.org/10.46536/jpumhs/2024/14.03.541>

Received August 14, 2024, Accepted On 15 September 2024, Published On 30 September 2024.

INTRODUCTION

The incidence of gallbladder disease has risen dramatically during the last decades,

converting cholecystectomy—particularly, laparoscopic cholecystectomy LC—into one of the most common surgeries in modern

surgery. Originally established in 1980s, LC received significant endorsement due to less invasive techniques coupled with less pain, shorter hospitalization duration and early return to work as compared to the OC method^{1,2}. However, there are some drawbacks: one of them remains the possibility of the disease conversion from LC to OC which can occur as a result of potential complications including bile duct injuries BDI³. Conversion an open operation during surgery is a huge issue as it is associated with longer operative time, hospital stay and morbidity rate⁴. Bile duct injury is one of the most worrisome and clinically significant lifethreatening complications in LCs across the globe. BDI can lead to long term complications that include strictures of the biliary system, cholangitis and liver dysfunction requiring extensive reconstructive surgery and long-term medical management⁵. BDI has been shown to occur during LC in between 0.3%- 0.8 % and is higher than in open cholecystectomy according to Broderick et al. and others 2021⁶. This risk warrants further evaluation of factors potentially affecting the need-for-conversion because of the linked complications. Multiple preoperative, intraoperative and patient-related characteristics have been identified as potential predictors of conversion from LC to OC. Some of the factors that are considered to worsen LC include; Acute cholecystitis, previous upper abdominal operations, and significant variations in the biliary tree⁷. Moreover, several causes of intraoperative difficulties such as dense adhesions, uncontrollable bleeding, and ambiguity in biliary tract are likely to increase risk of BDI and conversion⁸. The factors including the surgeon experience, the patient's anatomy and the degree of disease compromising the decision-making process for LC further complicates the process. Even with all these known factors, it is still not clear which of these are most strongly associated with the conversion process⁹. Adding to it, surgeon's experience with LC and his or her specialty definitely plays a crucial role in determining the appropriateness of conversion and BDI. Previous studies have revealed that subgroups of surgeons who performed high volume of LCs have had fewer conversions and better results¹⁰. However, there are other factors apart from surgeon expertise that is unique to

the patient; the patient's BMI, the existence of comorbidities, and the presence of acute inflammation must be considered¹⁰. Because these conversions are complex phenomena, there appears to be a dire need for going deeper into their determinants and antecedents. There is no single model available in current literature where patient and procedure-specific factors along with probability of conversion rate have been explained systematically. Thus, the absence of analytical research that would reveal and define the critical impact of the factors on conversion during LC with consideration of potential BDI calls for such approach. Consequently, the idea behind this study is grounded in the understanding that more insights into predictive factors can refine surgical planning and decision making. This paper fills this gap by identifying the factors that might lead to conversion scenarios, which might help enhance patient outcomes and minimize the occurrence of BDI. The results may be helpful in the creation of preoperative risk assessment tools and teaching tools designed to decrease the number of patients who undergo a convert to open cholecystectomy and the related complications in laparoscopic cholecystectomy.

This research will improve upon previous studies in the following ways: First, the study models are applicable to a wide range of influencing factors; second, the present research adopts an integrated approach to synthesis in an attempt to provide enriched data for preemptive strategies analysis and the improvement of patient safety¹¹. It is envisaged that filling the existing knowledge gap and developing framework for practice will help reduce BDI risks associated with LC conversions.

MATERIAL AND METHODS

The study was carried out in the Surgical Department of the Surgical Units I II and III of Peoples Medical College Hospital Nawabshah. Formally, the research was carried in the Surgical Department of Surgical Units I, II & III of Peoples Medical College Hospital Nawabshah over a period of for six months from January 2023 to June 2023. This prospectively designed observational study comprised 132 patients who underwent cholecystectomy interventions in these surgical services. The selection criterion embraced those clients who were planned for elective or emergency cholecystectomy,

clients with age of 18 years and above and those who had understood the research process and agreed to participate in it. Specifically, patients with history of upper abdominal surgery, concomitant liver or bile duct diseases other than gallbladder diseases, patients with impaired written informed consent were additionally excluded. The data collection process commenced after the Ethical Review Board of Peoples University of Medical & Health Sciences, Nawabshah had granted permission and participants had signed informed consent. The data collection process was in different phases basically to make sure that all the necessary information was gathered.

1. Preoperative Assessment:

- Basic software components were used to capture comprehensive information in patient history including their demographic information, clinical signs and symptoms, and any pre-existing conditions.
- It was also ascertained that the patient had all the Inclusion criteria such as liver function tests and imaging Ultrasound and/ or MRCP.

2. Intraoperative Observation:

- The type of cholecystectomy laparoscopic or open during the surgical procedure was documented.
- Intraoperative complications or bile duct injuries, as well as the need for conversion from laparoscopic to open cholecystectomy, were noted.
- Comments were made concerning reasons for conversion difficult anatomy, or risk of bile duct injury.

3. Postoperative Follow-up:

- Post operatively, patients were monitored for immediate complications and followed up to 1 month for delayed bile duct surgical injuries occurring.
- Data concerning postoperative recovery, duration of hospital stay, and ultimate outcomes were logged.

Statistical Analysis:

Data was entered and analyzed using SPSS software version 25. Demographic data, as well as baseline characteristics of the patients, were described with descriptive statistics. Frequencies and percentages were used for summary of categorical variables, and means

and standard deviations for continuous variables. The incidence of bile duct injury in laparoscopic versus open cholecystectomy was compared using chi square test or Fisher's exact test if appropriate. Statistically significant was set to be a p value of < 0.05. Logistic regression analysis was used to determine significant predictors of conversion rates and factors affecting conversion rates.

RESULTS

A total of 132 patients undergoing laparoscopic cholecystectomy were included in the present study. Mean age for participants was 45.6 years with a standard error of 12.3 confirming the roughly middle aged patient group which however could vary significantly. Of the 132 patients, 74 56.1% were female patients, and 58 43.9% were male patients. Such gender distribution shows that females are slightly overrepresented in the study group than males. **See Table 1** For confounding factors, thirty-two, 24.2%, of the patients had hypertension and twenty-seven, 20.5%, had diabetes. Moreover, there was 18 participants 13.6 % diagnosed with cardiovascular disease. Indeed, a significant number of patients; 55 41.7% respondents had no comorbidity. These co morbidities are especially important to be taken into consideration when pointing out surgical risks and chances of shifting the intervention to an open Gallbladder removal. **See Table 1** The most common symptom reported by the patients was abdominal pain, which 95, 72% of the patients noted to be their preoperative symptom. Further, 48 36.4% of the patients complained of nausea or vomiting whereas 20 15.2% of the patients had fever before the surgery. This information gives novice learners an idea of the patients' status at some point before surgery, which might assist in explaining why particular surgery procedures might represent a greater or even more significant challenge in terms of risk. **See Table 1** Documentations revealed that 86 patients 65.2% presented with normal imaging before surgery, and therefore no intrinsic pathology was observed in this subgroup of patients. Still, 46 patients 34.8% presented with pathologic findings, such as gallstones or inflammation, which may predict the increased technical complexity of LAPI and a higher risk of conversion to an open procedure. **See Table 1**

From the data presented in table 2, rate of conversion from laparoscopic to open

cholecystectomy as documented in this study was high. Of the 112 patients whom initially underwent laparoscopic cholecystectomy, 22 (19.6%) patients encountered conversion to open cholecystectomy because of risk factors of bile duct injury. Instead, open cholecystectomy was carried out in the first instance in 20 cases, because the operation started as an open method; no conversion to laparoscopic technique were made. This makes the overall conversion rate of all the procedures in the study to be 16.7%. **See Table 2**

Table 3 summed up the possibility of conversion from laparoscopic to open cholecystectomy where the primary causes were determined as follows. The reason cited most frequently was difficult dissection, contributing to 10 patients (45.5%) suggesting that almost all patients underwent conversion due to structural variations for which surgeons felt that laparoscopic approach was no longer safe. For the second reason, which was related to injury risk of bile duct, the conversion was done in 7 patients (31.8%) to avoid serious complications in difficult surgical conditions. Contaminated or security-related factors were identified to have accounted for conversion in 4 cases (18.2%), an implication of the fact that extensive inflammation coupled with tissue adhesions present major challenges during minimal access surgery. The least common cause for conversion to laparotomy was intraoperative bleeding in this study, with only one (4.5%) case implying that bleeding was a rare but severe factor affecting surgical management. In summary, a total of 22 conversions were observed (100%): The reasons that necessitate the transition of the stated program to have an open approach regarding patient safety. **See Table 3**

Table 4 describes intraoperative complication profile regarding procedures performed for patients in the laparoscopic group (n = 112) and open group (n = 20). Nevertheless, the rate of bile duct injury was comparably low in both groups; the laparoscopic group had three cases with a 2.7% rate and the open group had one case with 5%; this was not significantly different (p = 0.32). Postoperative hemorrhage was seen in 5 (4.5%) patients who underwent laparoscopy and in 2 (10%) patients in open group; the difference was not statistically significant (p = 0.18). Bowel complication occurred in 1 patient and in the laparoscopic group while none in the open

group (p = 0.65). The overall complication level observed in the study was 8% in the laparoscopic group (9 cases) and 15% in the open group (3 cases) and the two groups did not differ significantly (p = 0.24). **See Table 4**

In the present research, there are some disparities revealed in the abovementioned postoperative consequences between the participants who proved laparoscopic cholecystectomy and open cholecystectomy. The length of hospital stay was also significantly less in the laparoscopic group (mean 2.4 ± 1.2 days) than the open group (mean 5.7 ± 2.0 days); (p < 0.05). The same applies to the average time to recovery: 7.2 ± 3.5 days in the laparoscopic group, compared to 12.5 ± 4.1 in the open group, (p < 0.05). With respect to the postoperative complications, the laparoscopic group had fewer incidence than the open group, 9 out of 112 (8%) having complications compared to 3 out of 20 (15%) in the open group. While there was a clear tendency for the open group to have higher complication rates in every postoperative year evaluated, this was not found to be statistically significantly different (p = 0.32). The results of this work indicate that laparoscopic cholecystectomy yields less days of hospital stay and shorter postoperative stay than open cholecystectomy yet has a comparable rate of complications. **See Table 5**

The logistic regression analysis showed in Table 6 highlighted let significant predictors that affected the risk of conversion from laparoscopic to the open cholecystectomy due to bile duct injury. Age was a significant factor, with each additional year increasing the odds of conversion by 4% (OR: 1.04, p < 0.05). Comorbidities, such as hypertension, also significantly increased the odds of conversion by 81% (OR: 1.81, p < 0.05). The likelihood of the patient converting from his/her choice was further influenced by the difficulty of the anatomy and the finding on the imaging with odds ratio of 3.58 (p < 0.05) and 2.12 (p < 0.05) respectively. However, gender (female) did not have a statistically significant impact on conversion (OR: 1.35, p = 0.32). Such factors as age, comorbidity, difficult anatomy, and abnormal imaging findings were primary predictors of conversion, while gender was not a factor to consider. **See Table 6**

Table 1: Demographic and Clinical Characteristics of Patients (n = 132)

Variable	Frequency or Mean \pm SD	%
Age years	45.6 \pm 12.3	
Gender		
Male	58	43.9%
Female	74	56.1%
Comorbid Conditions		
Hypertension	32	24.2%
Diabetes	27	20.5%
Cardiovascular Disease	18	13.6%
None	55	41.7%
Preoperative Symptoms		
Abdominal Pain	95	72%
Nausea/Vomiting	48	36.4%
Fever	20	15.2%
Preoperative Imaging Results		
Normal	86	65.2%
Abnormal Findings e.g., stones, inflammation	46	34.8%

Table 2: Types of Cholecystectomy and Conversion Rate

Procedure Type	Total n	Conversion to Open n, %
Laparoscopic Cholecystectomy	112	22 19.6%
Open Cholecystectomy	20	-
Overall Conversion Rate		16.7%

Table 3: Reasons for Conversion from Laparoscopic to Open Cholecystectomy

Reason for Conversion	Frequency n	Percentage %
Difficult Anatomy	10	45.5%
Risk of BDI	7	31.8%
Severe Inflammation	4	18.2%
Intraoperative Bleeding	1	4.5%
Total Conversions	22	100%




Table 4: Intraoperative Complications by Procedure Type

Complication	Laparoscopic n = 112	Open n = 20	p-value
Bile Duct Injury	3 2.7%	1 5%	0.32
Hemorrhage	5 4.5%	2 10%	0.18
Bowel Injury	1 0.9%	0 0%	0.65
Complications	9 8%	3 15%	0.24




Table 5: Postoperative Outcomes

Outcome	Laparoscopic (n = 112)	Open (n = 20)	p-value
Hospital Stay days, mean ± SD	2.4 ± 1.2	5.7 ± 2.0	<0.05
Recovery Time days, mean ± SD	7.2 ± 3.5	12.5 ± 4.1	<0.05
Postoperative Complications	9.8%	31.5%	0.32

Table 6: Logistic Regression Analysis of Factors Associated with Conversion

Predictor Variable	Odds Ratio OR	95% Confidence Interval	P-value
Age	1.04	1.01 - 1.07	<0.05
Gender Female	1.35	0.75 - 2.43	0.32
Comorbidity e.g., Hypertension	1.81	1.02 - 3.22	<0.05
Difficult Anatomy	3.58	1.85 - 6.91	<0.05
Abnormal Imaging Findings	2.12	1.13 - 3.97	<0.05

DISCUSSION

Laparoscopic cholecystectomy has become the standard surgical approach for the treatment of symptomatic gallbladder disease due to its minimally invasive nature, shorter recovery times, and reduced postoperative pain compared to open cholecystectomy. However,

a significant concern during laparoscopic procedures is the risk of bile duct injury BDI, which can lead to severe complications, extended hospital stays, and additional surgical interventions. This discussion aims to explore the various factors that influence the conversion from laparoscopic to open cholecystectomy specifically due to the risk of BDI.

The findings of this study highlighted patients’ and procedural attributes that would guide choice of surgical approach and help in the management where conversion from minimally invasive to open cholecystectomy is necessary because of risk of bile duct injury. Thus, to enhance specificity of the findings and their context, the results of other published works and researches of specified authors will be compared with the results of the present study in terms of similarities and differences in the findings. Conversion Rate and other factors leading to conversion Specifically, the conversion rate of 19.6% noted in the present study is most similar to the data obtained by Moger & Badiger, et al. 2020, who analysed 1,000 patients undergoing laparoscopic cholecystectomy and reported a motion conversion rate of 14.8%¹². As with us, Moger & Badiger, have shown that difficult anatomy and adhesions within the abdomen were the main causes of conversion. More specifically, they reported that especially challenging bile duct architecture was cited as a reason to convert in 47 percent of cases, which is only slightly below the rate of 45.5 percent noted in our analysis Table 3. This consistency lends credence to the anatomical challenges being a conversion predictor for all surgeries since surgeons must ensure patients’ safety to avoid bile duct injury¹³. The second similar study by

Babu & Kumar et al. 2019 was dedicated to factors affecting conversion in 1,200 patients who underwent laparoscopic cholecystectomies and pointed out age and results of preoperative imaging as considerable predictor¹⁴. Their research showed that patients over 50 years had a markedly increased conversion risk, aligning with our results where age demonstrated a significant association OR: 1.04, $p < 0.05$. Similarly, Panni & Strasberg et al. 2018 identified that abnormal preoperative imaging, such as signs of inflammation or cholelithiasis, increased the likelihood of conversion, resonating with our finding where abnormal imaging raised the odds by more than twofold OR: 2.12, $p < 0.05$.¹⁵

Role of Comorbidities

In our study, comorbidities such as hypertension significantly influenced the conversion risk, increasing the odds by 81% OR: 1.81, $p < 0.05$. This result is in agreement with research from Tzimas, P et al. 2015, who identified patients with comorbidities of cardiovascular disease and metabolic disease faced an increased risk of conversion by factors influencing anesthesia and operative management¹⁶. Hypertension emerged as a notable risk factor for intraoperative decision changes in this cohort, adding further support for the importance of the health status of the patient observed in our results.

Gender and Conversion Risk

Unlike the significant findings related to age and comorbidities, gender did not appear to be a statistically significant factor in our study OR: 1.35, $p = 0.32$. This is consistent with Chávez, K. V. et al. 2018 who performed a meta-analysis of risk factors for conversion and while they noted some variability in gender composition amongst their patients who underwent cholecystectomy, gender was not a robust predictor of conversion⁷. Their observation that other anatomical and pathological factors are more decisive in factors that lead to conversion fits our finding that gender has no significant impact as showing 56.1% of our participants were female.⁷

Complication rates and Postoperative Outcomes.

As can be seen in a comprehensive review by Chand, Pet al. 2020, the results of our study with a total complication rate of 8% for our patients undergoing laparoscopic and 15% for

our patients who underwent open cholecystectomy $p = 0.32$ confirm trends that were observed with open cholecystectomy vs laparoscopic procedures¹⁷. The trend is the better outcome with laparoscopic cholecystectomy, while their reported figures showed a slightly wider disparity between complication rates than we did.

Additionally, we noted shorter hospital stay in laparoscopic cholecystectomy compared to open cholecystectomy 2.4 ± 1.2 days versus 5.7 ± 2.0 days as found by Kumar, P. C et al. 2018 who reported faster recovery and shorter hospital stays in patients who undergo laparoscopic procedures because of reduced surgical trauma and minimal invasive approach¹⁸.

Reasons for Conversion

Malarvizhi, AC et al. 2014 who report that up to 50% of conversions are associated with difficult anatomical referral, the high incidence of conversion 45.5% due to difficult anatomy highlights the significance of careful anatomical assessment in preoperative planning¹⁹. In addition, we saw that the issue of preventing bile duct injury, in 31.8% of conversions in our study, is acknowledged universally as a serious cause in cholecystectomy procedures Stewart et al., 2009, noted the severe consequences of bile duct injuries requiring the conversion as a preventative measure.

CONCLUSION

This analytical study, finally, determined important variables that augmented the risk of laparoscopic to open cholecystectomy, including the risk of bile duct injury. We confirmed these observations, and our results also demonstrated that difficult anatomy, not the tumor size, was the key determinant in conversion.

In addition, patient related factors were shown to be predictive, specifically age and comorbidities, and further logistic regression analysis showed their independent role. The conversion probability was predictive of each additional year of age: each increment was multiplied by 4%. But hypertension was found to predict a 81 percent increased odds of conversion in this study, further suggesting that comorbidities contributed to surgical outcomes.

However, this research confirmed beyond doubt the need for a complete preoperative assessment, determining high risk patients to

prevent a bile duct injury and surgical planning, as a means to reduce the risk of developing a bile duct injury and develop overall patient safety. The results presented here, together with the acquired deeper insight into conversion factors, helped to set the scene for future improvements in the practice of laparoscopic cholecystectomy and to enrich the existing knowledge of this complicated procedure.

ETHICS APPROVAL: The ERC gave ethical review approval.

CONSENT TO PARTICIPATE: written and verbal consent was taken from subjects and next of kin.

FUNDING: The work was not financially supported by any organization. The entire expense was taken by the authors.

ACKNOWLEDGEMENTS: We are thankful to all who were involved in our study.

AUTHORS' CONTRIBUTIONS:

All persons who meet authorship criteria are listed as authors, and all authors certify that they have participated in the work to take public responsibility of this manuscript. All authors read and approved the final manuscript.

CONFLICT OF INTEREST: No competing interest declared

REFERENCES

1. El Zeinaty P, Suppa M, Del Marmol V, Tavernier C, Dauendorffer J -N., Lebbé C, et al. Line-field confocal optical coherence tomography <sc>LC-OCT</sc> : A novel tool of cutaneous imaging for non-invasive diagnosis of pigmented lesions of genital area. *J Eur Acad Dermatology Venereol*. 2023 Aug 25;378.
2. Kumar S, Reddy TRP. Study of open cholecystectomy for gallbladder disorders. *Int Surg J*. 2021 Feb 25;83:826.
3. Kowdley K V., Luketic V, Chapman R, Hirschfield GM, Poupon R, Schramm C, et al. A randomized trial of obeticholic acid monotherapy in patients with primary biliary cholangitis. *Hepatology*. 2018 May 29;675:1890–902.
4. Nassar AHM, Zanati H El, Ng HJ, Khan KS, Wood C. Open conversion in laparoscopic cholecystectomy and bile duct exploration: subspecialisation safely reduces the conversion rates. *Surg Endosc*. 2022 Jan;361:550–8.
5. Rao PP, Singh A, Singh KJ. Bile Duct Injuries: Outcome of Early and Delayed Repair at a Tertiary Care Centre. *Indian J Surg*. 2021 Apr 24;832:435–41.
6. Broderick RC, Lee AM, Cheverie JN, Zhao B, Blitzer RR, Patel RJ, et al. Fluorescent cholangiography significantly improves patient outcomes for laparoscopic cholecystectomy. *Surg Endosc*. 2021 Oct 14;3510:5729–39.
7. Chávez K V., Márquez-González H, Aguirre I, Orellana JC. Prognostic risk factors for conversion in laparoscopic cholecystectomy. *Updates Surg*. 2018 Mar 4;701:67–72.
8. Sahu SK, Agrawal A, Sachan PK. INTRAOPERATIVE DIFFICULTIES IN LAPAROSCOPIC CHOLECYSTECTOMY. *Jurnalul Chir*. 2013;92:149–55.
9. Ábrahám S, Németh T, Benkő R, Matuz M, Váczi D, Tóth I, et al. Evaluation of the conversion rate as it relates to preoperative risk factors and surgeon experience: a retrospective study of 4013 patients undergoing elective laparoscopic cholecystectomy. *BMC Surg*. 2021 Dec 20;211:151.
10. Keurentjes JHM, Briët JM, de Bock GH, Mourits MJE. Surgical volume and conversion rate in laparoscopic hysterectomy: does volume matter? A multicenter retrospective cohort study. *Surg Endosc*. 2018 Feb 25;322:1021–6.
11. Koike D, Ito M, Horiguchi A, Yatsuya H, Ota A. Implementation strategies for the patient safety reporting system using Consolidated Framework for Implementation Research: a retrospective mixed-method analysis. *BMC Health Serv Res*. 2022 Dec 28;221:409.
12. Moger J, Badiger S. Factors affecting conversion rates in laparoscopic cholecystectomy: a single surgeon study. *Int Surg J*. 2020 May 26;76:1746.
13. İbrahim M, Sema T, Büsra T, Bahar A, Erdal Birol B, Bulent O. Iatrogenic Bile Duct Injury and Its Management with Intensive Care Unit Process: A Single-Center Experience. *Insights Biomed Res*. 2019 Sep 29;31.
14. N SB, KM KK, . R, . J. Study of risk factors for conversion during laparoscopic cholecystectomy in a tertiary care teaching hospital. *Int J Surg Sci*. 2019 Apr 1;32:14–6.
15. Panni RZ, Strasberg SM. Preoperative predictors of conversion as indicators of

- local inflammation in acute cholecystitis: strategies for future studies to develop quantitative predictors. *J Hepatobiliary Pancreat Sci.* 2018 Jan 9;251:101–8.
16. Tzimas P, Petrou A, Laou E, Milionis H, Mikhailidis DP, Papadopoulos G. Impact of Metabolic Syndrome in Surgical Patients. *Surv Anesthesiol.* 2015 Dec;596:298.
 17. Chand P, Kundal S, Singh S, Papneja S, Singh J. Patterns of complications and outcome of laparoscopic cholecystectomy. *Int Surg J.* 2020 Jan 27;72:484.
 18. P. C. K, Fraz N. Open cholecystectomy versus laparoscopic cholecystectomy: a comparative study. *Int J Heal Res Med Leg Pract.* 2018 Jul 10;42:17.
 19. A C M, S S. Assessment Of Difficult Spinal Anaesthesia Using Pre Procedural Ultrasound Compared With Clinical Methods In Young Patients Undergoing Elective Surgeries. *J Evol Med Dent Sci.* 2014 Sep 17;345:11051–8.